

## Vascular changes in the cycling and early pregnant uterus

Noura Massri, ... , Ripla Arora, Nataki C. Douglas

*JCI Insight.* 2023;8(11):e163422. <https://doi.org/10.1172/jci.insight.163422>.

### Review

Uterine vascular remodeling is intrinsic to the cycling and early pregnant endometrium. Maternal regulatory factors such as ovarian hormones, VEGF, angiopoietins, Notch, and uterine natural killer cells significantly mediate these vascular changes. In the absence of pregnancy, changes in uterine vessel morphology and function correlate with different stages of the human menstrual cycle. During early pregnancy, vascular remodeling in rodents and humans results in decreased uterine vascular resistance and increased vascular permeability necessary for pregnancy success. Aberrations in these adaptive vascular processes contribute to increased risk of infertility, abnormal fetal growth, and/or preeclampsia. This Review comprehensively summarizes uterine vascular remodeling in the human menstrual cycle, and in the peri- and post-implantation stages in rodent species (mice and rats).

**Find the latest version:**

<https://jci.me/163422/pdf>



# Vascular changes in the cycling and early pregnant uterus

Noura Massri,<sup>1,2</sup> Rachel Loia,<sup>3</sup> Jennifer L. Sones,<sup>4</sup> Ripla Arora,<sup>2,5</sup> and Nataki C. Douglas<sup>6,7</sup>

<sup>1</sup>Cell and Molecular Biology Graduate Program and <sup>2</sup>Institute for Quantitative Health Science and Engineering, Michigan State University, East Lansing, Michigan, USA. <sup>3</sup>School of Graduate Studies, Rutgers Biomedical and Health Sciences, Newark, New Jersey, USA. <sup>4</sup>Veterinary Clinical Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, Louisiana, USA. <sup>5</sup>Department of Obstetrics, Gynecology and Reproductive Biology, Michigan State University, East Lansing, Michigan, USA. <sup>6</sup>Department of Obstetrics, Gynecology and Reproductive Health and <sup>7</sup>Center for Immunity and Inflammation, Rutgers Biomedical and Health Sciences, Newark, New Jersey, USA.

**Uterine vascular remodeling is intrinsic to the cycling and early pregnant endometrium. Maternal regulatory factors such as ovarian hormones, VEGF, angiopoietins, Notch, and uterine natural killer cells significantly mediate these vascular changes. In the absence of pregnancy, changes in uterine vessel morphology and function correlate with different stages of the human menstrual cycle. During early pregnancy, vascular remodeling in rodents and humans results in decreased uterine vascular resistance and increased vascular permeability necessary for pregnancy success. Aberrations in these adaptive vascular processes contribute to increased risk of infertility, abnormal fetal growth, and/or preeclampsia. This Review comprehensively summarizes uterine vascular remodeling in the human menstrual cycle, and in the peri- and post-implantation stages in rodent species (mice and rats).**

## Introduction

Under physiologic conditions, most vessels in the body are highly stable once formed. Blood vessels in the mammalian uterus are an exception, with characteristic structural and functional changes in both the non-pregnant and pregnant states (1, 2). In humans, in the absence of pregnancy, ovarian-derived estrogen and progesterone mediate cyclic vascular remodeling in the endometrium (1–6). In early pregnancy in rodents and humans, uterine vascular remodeling supports embryo growth and development and subsequent placenta formation (7, 8). Abnormalities in the uterine vasculature increase the risk of infertility, miscarriage, and pregnancy complications such as fetal growth restriction and preeclampsia. Thus, the normal physiologic vascular changes that occur during pre-pregnancy and early pregnancy are likely evolutionarily selected to support pregnancy success. Due to ease of visualization and relationship to the placenta, most documented changes in uterine vasculature have extensively focused on post-placental stages of pregnancy, and a discussion of uterine vasculature prior to and in early stages of pregnancy is missing (7, 9). Here, we describe uterine vasculature anatomy in both humans and rodents and factors that regulate vascular remodeling during the menstrual cycle for humans and the pre-placentation stages of rodent pregnancy. Uterine spiral arterioles undergo clinically significant adaptations during rodent and human pregnancy and have been more extensively investigated than uterine veins; these vessels and their derivatives are the focus of this Review.

## Human uterus anatomy and the menstrual cycle

The human uterus is composed of the myometrium, a smooth muscle layer; the endometrium, an inner mucosal lining surrounding mesenchymal stroma; immune cells; and an extensive vascular network (10). The endometrium is further divided into the deep stratum basalis, adjacent to the myometrium, and the superficial stratum functionalis. The normal human menstrual cycle lasts 21 to 35 days and is divided into the estrogen-dominated proliferative phase, which has marked endometrial layer thickening, and the progesterone-dominated secretory phase (Figure 1A) (10). Ovulation, with oocyte release and corpus luteum formation, marks the beginning of the secretory phase and progesterone-mediated endometrial differentiation in preparation for embryo implantation. In the absence of pregnancy, the endometrium and associated vasculature are shed in response to the demise of the ovarian corpus luteum and withdrawal of ovarian hormones.

**Authorship note:** RA and NCD are co-senior authors.

**Conflict of interest:** NCD is chair of the Modern Fertility Medical Advisory Board.

**Copyright:** © 2023, Massri et al. This is an open access article published under the terms of the Creative Commons Attribution 4.0 International License.

**Reference information:** *JCI Insight*. 2023;8(11):e163422. <https://doi.org/10.1172/jci.insight.163422>.

Thus, with each cycle, the functionalis layer responds to estrogens and progesterone, is shed with menstruation, and then regenerates from the basalis layer (10, 11). Endometrial hypoxia and vasoconstriction of spiral arterioles are thought to limit the amount of blood lost during menstruation (12).

### Types of blood vessels in the human endometrium

Uterine vasculature comprising both blood and lymphatic vessels has been consistently observed in the human myometrium (13). In contrast, in the endometrium, there are conflicting reports about the location of lymphatic vessels, while blood vessels have been well characterized (13–16). Blood vessels localize to three regions of the human endometrium: the subepithelial zone, functionalis layer, and basalis layer. The subepithelial zone extends to less than 200  $\mu\text{m}$  below the surface epithelium, the basalis layer extends to less than 300  $\mu\text{m}$  above the myometrium, and the functionalis is between (17). Uterine arteries are located within the myometrium and serve as the major blood supply to the uterus, branching into the arcuate and then the radial arteries. At the myometrial-endometrial junction, the radial arteries split to form basal arteries, which supply the basalis layer, and spiral arterioles, which supply the functionalis layer. Arteries in the basalis consist of endothelial cells associated with vascular smooth muscle cells (VSMCs) that express  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA),  $\gamma$ -SMA, and myosin heavy chain. As the vessels course toward the luminal epithelium, VSMC content is reduced. Capillaries branching off from the spiral arterioles are present throughout the stroma and around the glandular epithelium. Beneath the luminal epithelial lining of the endometrium, capillaries arising from the spiral arterioles contribute to the subepithelial capillary plexus (3, 18). Microvessels at the subepithelial luminal surface are devoid of mural cells, including VSMCs and pericytes, and consist only of endothelial cells.

### Factors regulating the human endometrial vasculature


Cyclic changes in the endometrial vasculature in response to ovarian steroid hormones are associated with angiogenic factor activation.

#### Steroid hormones

Changes in the endometrial vasculature during the menstrual cycle are thought to be directly regulated by estrogens and progesterone. Localized expression of estrogen and progesterone receptors in the functionalis layer has been described in endometrium from hysterectomy specimens. Estrogen receptor 1 (ESR1) and ESR2, previously described as estrogen receptors (ERs)  $\alpha$  and  $\beta$ , respectively, and progesterone receptor (PR) are expressed in VSMCs (19). ESR1 is the predominant ESR during the late follicular phase, and ESR2 is predominant in the late secretory phase, suggesting hormonal regulation of their expression (19, 20). Whereas both ESR and PR are expressed in isolated human endometrial endothelial cells (21), endothelial cell expression of ESR2 has been detected in both proliferative- and secretory-phase endometrium, whereas PR expression has been variably detected (20–22). Thus, vascular remodeling could be mediated by acutely hormone-sensitive hormone receptor-expressing endothelial cells and VSMCs.

#### Angiogenic factors

The endometrial expression of angiogenic factors, VEGF, angiopoietins (ANGPTs), FGF, and their receptors implicates involvement in regulating vascular changes in the proliferative and secretory phases (21, 23). VEGF is known to promote de novo blood vessel growth, angiogenesis, and vascular permeability via tyrosine kinase family receptors, including VEGFR1 (FLT1), VEGFR2 (KDR/FLK1), and VEGFR3 (FLT4) (24). In both the proliferative and secretory phases, VEGFA is expressed in neutrophils associated with the subepithelial capillary plexus and in microvessels in the functionalis layer (25). In the proliferative phase, VEGFA expression also localizes to glandular epithelial and stromal cells (23, 26), and VEGFR2 is found in endothelial cells (23). In the secretory phase, uterine natural killer (uNK) cells express high levels of the angiogenic factors VEGFC, placental growth factor (PGF), and VEGFR3. VEGFA expression is higher in the epithelium than in the stroma. In the secretory phase, endothelial VEGFR2 levels decline, whereas VEGFR1 levels increase (23, 26). In proliferative and secretory endometrium, ANGPT1 is predominantly expressed in stromal cells, while ANGPT2 and the angiopoietin receptor TEK receptor tyrosine kinase (TIE2) are highly expressed in endothelial cells (27). FGF1 and FGF2 are expressed in the epithelium and stroma, with higher expression in the epithelium throughout the cycle (23). Isolated human endometrial endothelial cells proliferate in response to VEGF or FGF2, and addition of  $17\beta$ -estradiol increases, while progesterone decreases, endothelial cell proliferation in response to these angiogenic growth factors (21).



**Figure 1. Comparison of the human menstrual cycle and rodent estrous cycle. (A)** The human menstrual cycle is divided into the proliferative (follicular) phase and the secretory (luteal) phase. Under the influence of ovarian estrogen (blue line) in the follicular phase, the endometrium regenerates, endothelial cells proliferate, and vascular permeability increases. Mid-cycle, a surge in luteinizing hormone results in follicle rupture, ovulation of the oocyte, and formation of the corpus luteum. In the secretory phase, progesterone (red dashed line) regulates stromal cell decidualization and increases endometrial edema in preparation for embryo implantation. In the absence of embryo implantation, the corpus luteum demises, and estrogen and progesterone levels fall. This is followed by menstruation and the start of another menstrual cycle. **(B)** The rodent estrous cycle is composed of four different phases: proestrus, estrus, metestrus, and diestrus. In the proestrus phase, which lasts between 21 and 32 hours, estrogen increases. The estrus phase and ovulation follow the proestrus phase. The estrus phase lasts 12–48 hours. The formation of the corpus luteum following ovulation marks the shift from an estrogen- to a progesterone-dominated stage. The metestrus and diestrus phases represent the secretory phases of the cycle. The metestrus phase lasts 8–24 hours, while the diestrus phase lasts 48–72 hours.

## Vascular remodeling in the human menstrual cycle

### Mechanisms of angiogenesis

Angiogenesis can occur via three mechanisms: sprouting, elongation, or intussusception, each requiring unique changes in endothelial cells (Table 1). Endothelial cell expression of  $\alpha_v\beta_3$  integrin, a marker of vascular sprouts in non-endometrial tissues (28), has been detected only inside existing vessels and not in any sprout-like vascular structures. This observation suggests that elongation and intussusception and not sprouting angiogenesis are the primary mechanisms of angiogenesis in the human endometrium (20). The following sections highlight the roles that VEGF signaling might play in the cyclic remodeling of the human endometrial vasculature.

### Angiogenesis during proliferative phase

In the proliferative phase, estradiol, the primary ovarian estrogen, promotes rapid endometrial growth and angiogenesis. Stereologic examination of vessel length and branch points showed that vascular length density in the subepithelial capillary plexus is greatest during the mid to late proliferative phase. These observations are consistent with vessel elongation as the major mechanism of endometrial angiogenesis in the estradiol-dominated proliferative phase (17). Fine spatial analysis revealed a greater percentage of proliferating endothelial cells in the basalis layer during the proliferative phase compared with the secretory phase. In contrast, the proportion of proliferating vessels in both the subepithelial capillary plexus and functionalis layer was relatively constant across the menstrual cycle (25). Estradiol increases endothelial cell proliferative response to VEGF and promotes elongation angiogenesis and vascular permeability (20). A significantly greater percentage of VEGF-expressing vessels and more proliferating vessels with VEGF-expressing cells were observed in proliferative compared with the secretory endometrium for all endothelial layers evaluated (subepithelial capillary plexus, functionalis, and basalis). VEGF displayed higher expression in

**Table 1. Definitions of terms related to uterine vasculature**

Vasculogenesis	De novo formation of a vascular network through the differentiation of precursor cells (angioblasts) into endothelial cells that are then assembled into a primitive capillary plexus.
Angiogenesis	The formation and/or growth of new vessels from preexisting blood vessels.
Vascular sprout	An endothelial cell with an extension that is often directed toward a VEGF-producing cell.
Sprouting angiogenesis	The type of angiogenesis that involves formation of vascular sprouts from preexisting vessels to create new vessels.
Elongation angiogenesis	A type of non-sprouting angiogenesis that involves elongation and dilation of existing uterine blood vessels.
Intussusceptive angiogenesis	The type of non-sprouting angiogenesis in which proliferation of endothelial cells inside a vessel produces a wide lumen that can be split, forming an intussusceptive pillar.
Basal arteries	Endometrial vessels that branch from the uterine radial artery and supply blood to the basalis layer of the human endometrium.
Spiral arterioles	Endometrial vessels that are in the functionalis layer of the non-pregnant cycling human endometrium. These arterioles undergo extensive remodeling during pregnancy and are the main vessels connecting the uterus and the placenta.
Utero-ovarian or arcuate arteries	The rodent uterine and ovarian arteries anastomose to form the utero-ovarian arteries.
Circumferential arterioles	These arterioles run dorsally and ventrally between the circular and longitudinal muscle layers of the rodent uterus, and they supply blood to the myometrium and the endometrium.
Vascular permeability	The ability of the blood vessel wall to allow for flow of substances in and out of the vessel. In early pregnancy, increased vasculature permeability can be demonstrated by intravenous injection of Evans blue dye. Evans blue dye binds to albumin and leaks from endometrial blood vessels at implantation sites. The dye forms visible blue bands in the uterine horns, and each band represents an implantation site.
Gestational day (GD)	Mice are set up to mate during late evening, and a mating event is identified by the presence of a vaginal copulation plug that is visible; the day the vaginal plug is observed is termed gestational day 0.5 (~12 hours after the mating event).
Estradiol/estrogen	Steroid hormone produced by the growing ovarian follicle.
Progesterone	Steroid hormone produced by the ovarian corpus luteum.
VEGF	VEGF, also known as vascular permeability factor, was first identified as an essential growth factor for vascular endothelial cells. VEGF binds to receptors including VEGF receptor 1 (VEGFR1), VEGFR2, VEGFR3, and neuropilin 1 (NRP1).
Angiopoietins (ANGPTs)	Glycosylated, secreted proteins that modulate angiogenesis by binding to their endothelial receptors TIE1 (tyrosine kinase with immunoglobulin-like and EGF-like domains 1) and TIE2 (TEK receptor tyrosine kinase).
Notch	A family of type 1 transmembrane proteins forming a highly conserved signaling pathway that governs cell fate decisions in many tissues. Notch proteins (Notch1, Notch2, Notch3, and Notch4) bind to ligands delta-like ligand 1 (DLL1), DLL3, and DLL4 and Jagged1 (Jag1) and Jag2 to activate the pathway.

Uterine natural killer (uNK) cells Lymphocytes implicated in regulation of vascular remodeling during early pregnancy in both humans and rodents.

the subepithelial capillary plexus compared with the functionalis and basalis layers. Finally, the observation of focal VEGF in marginating and adherent neutrophils associated with microvessels implicates neutrophils as the primary intravascular VEGF source for microvessels undergoing angiogenesis (25).

### Secretory-phase angiogenesis

Vascular development in the progesterone-dominated secretory phase is thought to occur via intussusceptive angiogenesis (3). Structurally, spiral arterioles increase in size and tortuosity, and the subepithelial capillaries dilate during the secretory phase (29). Microvessel concentration is thought to significantly increase in the mid-secretory phase (29). After ovulation, VEGFR2 levels decline, whereas VEGFR1 levels increase. While VEGFR2 signaling in the proliferative phase promotes endothelial cell proliferation, VEGFR1 signaling is thought to increase endothelial cell migration in the secretory phase (23). Unlike endothelial cells, mural cells proliferate under the control of progesterone during the secretory phase and wrap around endothelial cells, regulating angiogenesis and coordinating the function of blood vessels. In the late secretory phase, perivascular mural cells form a thick layer and surround the spiral arterioles (26). uNK cells increase in number during the mid-secretory phase and are located close to the endometrial glands and spiral arterioles, suggesting a role in the vascular remodeling that occurs in this phase (30).

### Vascular breakdown during menses

At the end of the secretory phase, right before menstruation, there is a rapid regression of the endometrium that induces increased coiling of spiral arterioles. Menstruation is associated with changes in the

endometrial balance between matrix metalloproteinase (MMP) activators and tissue inhibitors of MMPs (TIMPs). Progesterone withdrawal seems to stimulate MMP activity, resulting in extracellular matrix (ECM) breakdown and bleeding. The transition from menses to the proliferative phase initiates with estradiol-independent repair of vascular beds during menstruation (29).

### The rodent estrous cycle

Like the human menstrual cycle, the rodent estrous cycle is orchestrated by ovarian-derived hormones that prepare the uterus for embryo implantation and pregnancy. In rodents, cyclic uterine remodeling occurs over 4–5 days (Figure 1B). Remodeling includes cell shape changes in the luminal and glandular epithelium and stromal cells, changes in uterine gland secretions, and changes in uterine lumen size and shape based on the volume of uterine luminal fluid (31). Although mice do not menstruate, there are mouse models of menstruation where endometrial shedding is induced after decidualization by withdrawal of progesterone (32). Despite these models, while a lot is known about the vascular remodeling events during the human menstrual cycle, little is known about vascular changes in the rodent estrous cycle.

### Types of blood vessels in the rodent uterus

Like in humans, in rodents, both blood and lymphatic vessels have been consistently described in the uterine myometrium, while the endometrium and decidua of pregnancy appear to be devoid of lymphatic vessels (33). The rodent uterus is radial but asymmetric with large uterine arteries present along the mesometrial pole (Figure 2A). The uterine arteries, formed from the anastomoses of the uterine and ovarian arteries, are referred to as utero-ovarian or arcuate arteries. These arteries supply blood to segmental arteries in the mesometrium that branch into circumferential arterioles that pass through the uterine smooth muscle layers and wrap around the uterus. Circumferential arterioles further branch into spiral arterioles that supply blood to the inner muscle layers, the endometrium, and subepithelial capillaries present in proximity to the luminal epithelium (Figure 2A) (4).

### Events in early rodent pregnancy


Mating events in rodents are easily documented by detection of a copulation plug, termed gestational day (GD) 0.5. After mating, fertilization takes place in the oviduct (the rodent fallopian tube) followed by a series of cell divisions to form the blastocyst-stage embryo. In the mouse, multiple blastocysts enter the uterus early on GD3.0, followed by embryo spacing and embryo attachment on GD4.0 (34). Embryos attach to the uterine lumen, forming an implantation chamber (35), followed by a rise in vessel permeability around the site of attachment at GD4.5 (36). The implantation chamber then invades into the underlying stroma as the luminal epithelium around the embryo undergoes cell death and the stromal cells undergo a mesenchymal-to-epithelial differentiation process termed decidualization (36, 37). Between GD5.5 and GD7.5, the embryo continues to develop in the decidua, and embryonic umbilical vasculature fuses with maternal circulation by GD8.5 to form the placenta by GD9.5 (36, 37). Rat gestation is about 2 days longer than that of the mouse, and all events described above are shifted accordingly (38).

Limited information is available on early human pregnancy owing to lack of noninvasive methodologies to study the uterine microenvironment. Therefore, rodent models have been extensively used to examine endometrial changes required for successful embryo implantation and pregnancy progression. In the following sections, we describe several such rodent models and highlight human studies where applicable. Tables 2–4 summarize additional factors not discussed in the text.

### Vascular remodeling in rodent pregnancy

#### Vascular modifications in rodents during peri-implantation (GD0.5–GD4.5)

In rat pregnancy, endothelial cell proliferation is increased in the entire endometrium in the first few days following ovulation (39, 40). In rats and mice, as embryos enter the uterus, there is a substantial increase in the diameter of subepithelial capillaries in the endometrium (Figure 2B) (5, 41). Embryos attach to the anti-mesometrial pole and subepithelial capillaries in proximity to the implantation region dilate, leading to increased vessel permeability following embryo attachment (36, 42, 43). It is thought that the loss of vascular endothelial cadherin (VE-cadherin) facilitates this increase in permeability (44). In rodents, intravenous blue dye injection prior to euthanasia aids in the identification of embryo implantation sites owing to



**Figure 2. Vascular remodeling from pre-implantation through decidualization.** (A and B) Longitudinal cross sections of the GD3.5 (A) and the GD4.5 (B) mouse uterus illustrating arteries (red), veins (blue), and the uterine lumen (outlined in gray). The utero-ovarian artery (UA) and vein (UV) give rise to the segmental arteries and veins, respectively (SA and SV). The segmental arteries divide into circumferential arterioles (CA) that then branch to form the spiral arterioles (SpA). Subepithelial capillary plexus (SCP) arise from the spiral arterioles and supply blood to the endometrium. Following embryo attachment at GD4.5 (B), the vessels remodel and dilate in proximity to the implantation region. (C and D) Transverse cross sections at GD5.5 (C) and GD7.5 (D). At GD5.5 (C) the newly formed decidual capillaries are readily apparent. By GD7.5 (D), the vasculature surrounding the implantation chamber can be divided into different regions – the central region surrounding the embryo (CTR), the mesometrial region (MR), and the anti-mesometrial region (AMR) – each with a unique composition of vessels. The MR is composed of spiral arterioles and capillaries. The spiral arterioles contain endothelial cells and mural cells (pericytes and VSMCs). The decidual capillaries in the AMR are smaller vessels composed of endothelial cells associated with pericytes, while the decidual capillaries in the CTR contain very few pericytes. AM, anti-mesometrial pole; Cx, cervix; E, embryo; M, mesometrial pole; Myo, myometrium; O, ovary; UF, uterine fold.

vascular leakage through permeable vessels (36, 43). Increased vascular permeability at implantation sites across several mammalian species, including sheep and pigs, underscores the importance of this vascular change for successful implantation and later pregnancy events (45–47).

### Factors contributing to increased vascular permeability (GD0.5–GD4.5)

During pre-implantation stages, the factors that contribute to increased endothelial cell proliferation and the exact timing of this process are not well described (39, 40). In contrast, increased vascular permeability around implantation sites has been heavily studied, and several factors have been implicated in this process (Table 2), as discussed below.

**Steroid hormones.** In rodents, ovulatory estrogen induces epithelial proliferation at GD1–GD2, and a small rise in estrogen levels along with high levels of progesterone supports implantation by GD4.0 (48, 49). Experiments conducted in rodents to assess estrogen effects on the vasculature are typically performed with high estrogen levels, which mimic pre-ovulation conditions. Treatment of ovariectomized rats with estradiol or the estrogen agonist tamoxifen induced VEGF transcription (24, 50). Similarly, 17 $\beta$ -estradiol treatment of ovariectomized mice induced VEGF and VEGFR2 expression in the stroma within 6 hours of treatment (51). Further, a single injection of high-dose 17 $\beta$ -estradiol stimulated VEGF expression in the stroma within 2 hours after treatment (52). These studies imply a role for ovulatory estrogen in promoting expression of VEGF family genes. Like estrogen, progesterone alone induced stromal VEGFR2 expression in ovariectomized mice, although expression occurred within 12 hours of treatment. Thus, increase in serum progesterone levels may mediate the changes in uterine VEGF signaling that accompany embryo implantation (24, 51, 53, 54).

**VEGF signaling.** The VEGF receptors, VEGFR1, VEGFR2, VEGFR3, and neuropilin 1 (NRP1) are expressed in the peri-implantation mouse uterus (24, 55), and endothelial VEGF signaling in the peri-implantation period is required for embryonic growth and pregnancy progression. VEGF164, the

**Table 2. GD0.5–GD4.5: factors that regulate angiogenesis and vascular remodeling**

Gene	Expression		Inhibition/overexpression model		Refs.	
	Method	Developmental stages	Cell type and/or location	Method		Phenotype
<i>Ptgs2</i>	mRNA (ISH, NB) Protein (IHC)	GD3–GD4	Luminal epithelium; subepithelial stroma	Knockout mice	NA	61, 62, 65, 123
<i>Vegf</i>	mRNA (ISH, NB)	GD1–GD4	Luminal epithelium; subepithelial stroma; stromal bed	Timed administration of VEGF-blocking Ab (GD2)	Decreased vascular permeability at implantation sites	24, 57, 68, 79
<i>Vegfr2</i>	Protein (IHC, IF)	GD3.5	Endometrium; myometrium	NA	NA	68

IF, immunofluorescence; IHC, immunohistochemistry; ISH, in situ hybridization; NB, Northern blot.

predominant VEGF isoform in the mouse uterus, is expressed in epithelial cells from GD0 to GD3 and in subepithelial stroma on GD2 and GD3. Following embryo implantation, VEGF164 is localized to the luminal epithelium and stromal cells surrounding the blastocyst (24). Interestingly, VEGF expression was elevated around embryo sites in early pregnancy but not during pseudopregnancy, suggesting that the presence of an embryo is necessary to induce expression between GD4.5 and GD6.5 (56). Systemic administration of a VEGF-blocking antibody in early rat pregnancy (GD2) lowered vascular permeability, reducing the number of implantation sites at GD4 without significant effect on microvascular density or endothelial cell proliferation (57). Thus, VEGF signaling plays a crucial role in pre-implantation vascular remodeling events associated with embryo attachment (57).

**Prostaglandins.** Prostaglandin (PG) synthesis is necessary for the increased vascular permeability and uterine angiogenesis associated with early pregnancy. Prostaglandin synthase 2–derived (PTGS2-derived) PGs (PGE and PGI<sub>2</sub>) are elevated in implantation sites of several mammalian species (58–63). In rodents, depletion of PG function during pre-implantation is associated with abnormal post-implantation events, including implantation failure, decreased vascular permeability, and/or impaired decidualization, resulting in embryo growth restriction between GD5 and GD7 (45–47, 60, 62, 64–66). *Ptgs2*<sup>-/-</sup> mice also show defects in implantation and vascular permeability. However, daily intraperitoneal administration of the stable PGI<sub>2</sub> analog carbaprostacyclin to *Ptgs2*<sup>-/-</sup> mice starting at GD3.75 improves implantation and restores VEGF and VEGFR2 expression, along with blood vessel numbers (61, 62, 65), suggesting that PG-depletion phenotypes are partly associated with disrupted VEGF signaling. Despite the association of both PG and VEGF signaling with increased vascular permeability in early pregnancy, the impact of PGs and VEGFs on vessel structure around the site of embryo attachment is not understood.

#### Vascular modifications in rodents during decidualization (GD5.0–GD8.5)

From GD4.5 to GD7.5, decidual angiogenesis creates a rich, new capillary network (Figure 2C) (6). At GD5.5, the decidual transcriptome shows upregulation of endothelial cell-associated genes (67), with peak endothelial cell proliferation at GD6.5 (68). Thus, it is not surprising that differences in the decidual vasculature are readily apparent by GD7.5. The mesometrial region (MR) contains spiral arterioles covered with VSMCs and pericytes. The central region (CTR) contains capillaries with little or no pericyte coverage, while the capillaries in the anti-MR (AMR) are closely associated with pericytes (Figure 2D) (6, 69). Comparison of the GD7.5 vasculature to the estrous-stage non-pregnant vasculature revealed a 14.9-fold increase in the vascular sprouts in the CTR and a 7.4-fold increase in the vascular sprouts in the AMR. During early pregnancy, the intussusceptive blood vessels also increase 2.8-fold and 2.4-fold in the CTR and AMR, respectively. Thus, both sprouting and intussusceptive angiogenesis contribute to formation and remodeling of decidual vessels in the mouse.

#### Factors contributing to increased vascular permeability and endothelial cell proliferation (GD5.0–GD8.5)

Post-implantation decidual angiogenesis and vascular remodeling, which includes the formation of new vascular networks, are regulated by key angiogenic signaling pathways (Table 3), including VEGF, ANGPT, and Notch (6, 67, 70, 71), and immune cells (72, 73), such as uNK cells (6, 74, 75), as discussed below.



*Steroid hormones.* Similar to the pre-implantation/implantation stages, both estrogen and progesterone influence angiogenic factor expression in the post-implantation uterus. De novo synthesis of aromatase-derived estrogen from the stromal cells promotes angiogenesis during decidualization. Treatment with the aromatase inhibitor letrozole reduced expression of the endothelial cell-specific marker CD31 in the stromal compartment. Further, letrozole promoted downregulation of angiogenesis-related factors, such as endothelial PAS domain protein 1 (EPAS1), ANGPT2, and ANGPT4. These observations indicate that local estrogen production from the stromal cells induces secretion of angiogenic factors that contribute to endothelial cell network formation during decidualization (76).

Post-implantation treatment with a PR antagonist, RU486, decreased VEGF (6) and *Angpt2* expression at GD6.5 and GD8.5 and impaired uterine blood vessel remodeling at GD8.5 (70, 77). Spatiotemporal colocalization of PR and VEGFA was observed in decidual stromal cells at GD4.5. Further, while progesterone treatment at GD4.5 and GD5.5 increased blood vessel density in the CTR and AMR and increased vascular sinus fold numbers at GD6.5 in control mice, this progesterone-induced increase was not observed in VEGFA-deficient mice. Thus, progesterone-mediated changes in vascular remodeling primarily occur through VEGF and ANGPT2 signaling (6, 70).

*Angiogenic factors.* After implantation, *Vegf* mRNA is expressed in decidual cells in the MR and AMR of implantation sites from GD5 to GD7 (24, 55). *Angpt2*, *Angpt3*, and *Tie2* are expressed in the endometrium starting after implantation at GD5 and reach maximum expression after decidualization at GD8.5 (61, 70). Another ANGPT family member, ANGPT4, is expressed in endometrial fibroblasts and endothelial cell populations during decidualization (78).

VEGF signaling via VEGFR1, VEGFR2, and VEGFR3 contributes to post-implantation vascular remodeling events (24, 55, 56, 68). Intraperitoneal administration of a VEGFR2-blocking antibody at GD3.75 reduced uterine vasculature at GD5.5, with subsequent embryo loss by GD10.5. VEGFR2 blockade in a pseudopregnant mouse model with oil stimulus also decreased uterine vasculature at GD6.5 (68), suggesting that a physical stimulus is sufficient to induce VEGF-dependent vascular changes. In contrast, although pre-conception and early pregnancy administration of VEGFR1- or VEGFR3-blocking antibodies reduced vascular density at GD5.5, embryo growth and pregnancy outcomes were not compromised. The phenotypes observed with VEGFA signaling blockade and conditional loss of one copy of *Vegfa* suggest that VEGF/VEGFR2 signaling promotes an increase in vascular density, enlargement of vascular sinus folds, and endothelial cell proliferation in the CTR and MR on GD4.5–GD8.5 (6). Together, these data suggest that VEGF/VEGFR2 is the only decidual VEGF signaling pathway essential for progression of pregnancy prior to placenta formation (6, 68, 79).

Studies in mice and women implicate aberrant VEGF expression in the decidua as the cause of later-stage pregnancy complications. The pregnant blood pressure high/5 (BPH/5) mouse is a superimposed preeclampsia model that exhibits placental defects, including shallow trophoblast invasion, inadequate spiral artery remodeling, and compromised fetal growth (80, 81). The decreased decidual vasculature observed in BPH/5 pregnancies at GD7.5 is associated with altered VEGF164, *Vegfr1*, *Vegfr2*, and *Plgf* expression (80). In the decidua of women with a history of recurrent pregnancy loss (RPL), VEGF mRNA and protein levels were significantly decreased in comparison with those without a history of RPL, who were undergoing voluntary termination of pregnancy (82). This observation suggests a link between disrupted VEGF signaling in early pregnancy and increased risk of miscarriage. Together, both mouse and human studies highlight the importance of VEGF signaling in decidual angiogenesis, placentation, and pregnancy success.

*Notch signaling.* Like VEGF proteins, the distinct expression pattern of Notch proteins and ligands in the peri-implantation uterus reflects unique roles for Notch signaling in decidual angiogenesis and vascular remodeling (33, 69). Blockade of the Notch ligand delta-like ligand 4 (DLL4) at GD4.5 altered vascular tip and stalk cell identity, leading to excessive decidual angiogenesis and compromised embryonic growth by GD9.5 (71). Endothelial-specific deletion of the Notch regulator ADAM10 resulted in abnormal vascular patterning, impaired decidualization, and pregnancy loss by GD6.5 (67). In this model, there was increased diameter of blood vessels, a characteristic honeycomb structure of vessels near the implantation site, and increased endothelial cells in the AMR at GD5.5. In contrast, endothelial cell-specific loss of the Notch ligand Jagged1 at GD4.5 resulted in aberrant *Dll4* expression and increased Notch signaling without a change in the decidual vasculature or interruption of pregnancy at GD7.5 (69). These studies suggest a role for Notch signaling in decidual angiogenesis.

**Table 3. GD5.0–GD8.5: factors that regulate angiogenesis and vascular remodeling**

Gene	Expression		Inhibition/overexpression model		Refs.	
	Method	Developmental stages	Cell type and/or location	Model		Phenotype
<i>Adam10</i>	Protein (IF)	GD5.0–GD6.5	Endothelial cells	Endothelial-specific knockout	Upregulation of endothelial cell-associated genes; abnormal vascular patterns; large vessels and increased CD31 <sup>+</sup> endothelial cells in the implantation sites; impaired decidualization at GD5.5 and embryo resorption at GD6.5	67
<i>Angpt2</i>	mRNA (RT-qPCR) Protein (IHC, IF, WB)	GD5.0–GD7.0	Decidualizing stromal cells at the mesometrial and anti-mesometrial poles	NA	NA	70
<i>Angpt1</i>	mRNA (ISH)	GD5.0–GD6.0	Secondary decidual cells and undifferentiated stromal cells along the myometrium	NA	NA	61
<i>Angptl4</i>	mRNA (ISH, RT-qPCR)	GD6.5–GD8.5	Stromal and endothelial cells	NA	NA	78
<i>Cx43</i>	Protein (IHC)	GD4–GD6	Stromal cells	Uterine-specific knockout	Impaired angiogenesis in stromal cells at GD6.0 and GD7.0; reduced <i>Angpt2</i> and <i>Angpt4</i> expression in endothelial cells at GD7.0	101
<i>Cx43</i>	Protein (IF, WB)	GD5.5–GD7.5	Stromal and decidual cells	Mice with dominant loss of function	Irregular organization of vascular sinusoids in the decidua at GD6.5 and GD7.5; decreased uNK cells at the mesometrial pole; increased mRNA and protein expression of angiogenesis genes VEGF, VEGFR1, VEGFR2, and FGF2 at the mesometrial pole at GD6.5	99
<i>Cx43</i>	Protein (WB)	NA	Human endometrial stromal cells	Pharmacologic inhibition or shRNA knockdown	Decreased expression of VEGF mRNA and protein and impaired decidualization	103, 130
<i>Dll4</i>	Protein (IF)	GD7.5	Decidual capillary endothelial cells	Timed administration of Dll4-blocking Ab (GD4.5)	Increase in decidual vascular density, decidual cell proliferation, and apoptotic cells at GD6.5; decreased decidual vascular density, proliferative decidual cells, and apoptotic cells at GD9.5	69, 71
<i>Fgf2</i>	Protein (IHC, IF, WB)	GD6.0–GD8.0	Rat whole uterus (bulk tissue) Decidualizing stromal cells in the secondary decidual zone	NA	NA	131–133
<i>Ptgs2</i>	mRNA (RT-qPCR)	GD5–GD7	Luminal epithelium, subepithelial stroma, mesometrial pole	Knockout mice Pharmacologic inhibition (celecoxib and indomethacin)	Decidual angiogenesis defects; decrease in CD31 <sup>+</sup> endothelial cells; decreased VEGF164 and VEGFR2 expression at GD5 and GD7 Decreased vascular permeability	61, 62, 65, 43, 58, 60, 66, 134
<i>Vegf</i>	mRNA (ISH, NB) Protein (IHC, IF)	GD4.5–GD8.5	Decidualizing stromal cells at the mesometrial and anti-mesometrial poles Trophoblast giant cells	Timed administration of VEGF-blocking Ab (GD2) VEGF trap (blockade of VEGF and PGF) at GD4.5	Decreased vascular permeability at implantation sites Pruned vessel structures and reduced blood vessel densities in both the central region and the anti-mesometrial region by 35% and 53%, respectively; embryo resorption by GD6.5	6, 24, 68, 79
<i>Vegfr2</i>	Protein (IF)	GD5.0–GD7.5	Endothelial cells throughout the decidua and myometrium Adjacent to the embryo	Timed administration of VEGFR2-blocking Ab (GD3.75)	Decreased proliferation of endothelial cells GD5.5; decreased vascular density GD7.5; embryonic lethality by GD10.5	68
<i>Vegfr3</i>	Protein (IF)	GD5.0–GD7.5	Endothelial cells in the primary decidual zone	Timed administration of VEGFR3-blocking Ab (GD3.75)	Decreased proliferation of endothelial cells GD5.5; decreased vascular density GD7.5	68

IF, immunofluorescence; IHC, immunohistochemistry; ISH, in situ hybridization; NB, Northern blot; RT-qPCR, quantitative reverse transcription PCR; WB, Western blot.

*uNK cells.* uNK cells are immune cells that contribute to uterine vessel remodeling (83, 84). uNK cell precursors are present in the blood vessels and the stromal compartment in the pre-implantation mouse uterus (GD2–GD3). After implantation, uNK cells are abundant in the decidua and undergo notable maturation and proliferation from GD5.5 to GD7.5 (85). In addition, uNK cells produce soluble factors, including VEGF, PGF, and Notch ligand DLL1, that contribute to decidual angiogenesis and vascular remodeling during early pregnancy (73, 74, 85, 86). Between GD4.5 and GD6.5, uNK cells and VEGF/VEGFR3 signaling coordinately regulate enlargement and elongation of uterine vascular sinus folds in the MR (6). Even though uNK cells do not express estrogen and progesterone receptors in both mice (87–89) and humans (90), they are regulated by ovarian hormone signaling (72, 91). This is likely through steroid hormone regulation of stromal cells that surround the uNK cells (90). Consequently, ovarian stimulation (superovulation) with pregnant mare serum gonadotropin and human chorionic gonadotropin results in decreased uNK density near the MR of the decidua, deficient cytokine secretion, compromised placental angiogenesis, and low embryonic weight at GD8.5 (92).

Antibody-mediated depletion of uNK cells at GD6.5 and GD7.5 resulted in reduced blood vessel density and vascular sinus folding in the decidua CTR at GD8.5 (6, 75). There are conflicting data regarding the vascular phenotypes in uNK cell-deficient mouse models. IL-15 is a key cytokine for uNK cell maturation and is expressed in the decidual stromal cells (85). *Il15*<sup>-/-</sup> mice lack uNK cells and display expression of angiogenesis genes without abnormalities in vascular density or endothelial cell proliferation at GD7.5 (93). Conversely, a uNK cell-deficient model created by depletion of immune cells (*Rag2*<sup>-/-</sup> *Il2rg*<sup>-/-</sup>) exhibits delayed mesometrial angiogenesis and maturation of decidual vessels at GD6.5, which compromises fetal development at GD9.5 (75). This study suggests involvement of uNK cells in promoting vascular branching and vessel pruning following implantation (75). However, in addition to uNK cell depletion, *Rag2*<sup>-/-</sup> *Il2rg*<sup>-/-</sup> mice lack B and T cells (alymphoid). This may explain the differences in the two uNK cell-deficient models.

In humans, uNK cells are present in the endometrium and decidua of early pregnancy. These uNK cells express VEGFA and VEGFC, suggesting a role in decidual angiogenesis (26, 84). In vitro studies revealed that human uNK cells produce angiogenic factors (94) and that estrogen promotes uNK cell motility and uNK cell-mediated angiogenesis (95). After exposure to asoprisnil, a PR modulator, uNK cell numbers and IL-15 expression were significantly reduced in human secretory-phase endometrium. Impaired spiral artery remodeling, characterized by increased VMSC content and thickened arteriole walls, was also observed (96, 97). Increased numbers of uNK cells as well as increased vascular density and vascular wall thickness were observed in RPL compared with normal pregnancy (98).

*Gap junction protein connexin 43.* Connexin 43 (*Cx43*), also known as Gap junction alpha-1 protein (*Gja1*), is expressed in decidual stromal cells on GD4 and GD6. Mice with a dominant-negative loss of Cx43 function exhibited an increased and irregular pattern of decidual sinusoids at the mesometrial pole, decreased uNK cells at GD6.5 and GD7.5, upregulation of angiogenic factors such as VEGFA and FGF2, and fetal growth restriction (99). Global Cx43 loss results in postnatal death, preventing any analysis of fertility in the adult mouse (100). However, conditional deletion of *Cx43* in uterine stromal cells, epithelium, and the circular muscle via PR-Cre (*PR*<sup>Cre</sup>) reduced expression of ANGPT2 and ANGPT4 at GD6 and GD7. Uterine-specific Cx43-deficient mice also displayed reduced endothelial proliferation and a rudimentary vascular network on GD6 and GD7, resulting in poor decidualization and embryo development (101). These studies reveal the importance of intercellular communication via gap junctions in establishing the uterine vascular network that is necessary for decidual angiogenesis, embryo growth, and pregnancy success (82, 102, 103).

### Vascular remodeling in rodents during placentation (GD8.5–GD10.5)

Uterine vascular remodeling during placenta formation is mediated by decidual stromal cells, local uterine immune cells, including uNK cells, and embryo-derived trophoblast cells. When trophoblast cells in the trophectoderm layer of the blastocyst contact the endometrium, the mural trophectoderm differentiates into invasive trophoblast giant cells (TGCs), and the polar trophectoderm gives rise to the extraembryonic ectoderm and ectoplacental cone (EPC) (104). The EPC differentiates into several types of trophoblasts, including endovascular TGCs, interstitial glycogen trophoblast cells, and cytotrophoblasts, that serve unique functions in placental development in rodents (105). From GD8.5 to GD10.5, the major vascular changes are spiral arteriole remodeling and formation of the placenta.

**Spiral arteriole remodeling.** Spiral arteriole remodeling in rodents and women occurs via trophoblast-independent and trophoblast-dependent mechanisms. Trophoblast-independent, decidua-associated vascular remodeling involves endothelial vacuolation and VSMC swelling, leading to apoptosis. Trophoblast-dependent vascular remodeling involves the action of endovascular trophoblasts, the subtype of invasive trophoblasts that invades the arteriole wall, resulting in a chimeric vessel composed of both trophoblast and endothelial cells (Figure 3).

Interstitial and endovascular trophoblast invasion is required for placenta formation in rodents and women (106). In both species, interstitial trophoblasts degrade ECM proteins and invade the decidua to arrive at the spiral arteriole adventitial layer, whereas endovascular trophoblasts percolate through the vessel wall, restructure smooth muscle, and displace the endothelium (106). For rodent models of placenta formation, an important difference between rats and mice is the depth of trophoblast invasion. In rats, endovascular and interstitial trophoblasts invade beyond the decidua into the mesometrial triangle (or metrial gland) of the myometrium, while in mice, endovascular trophoblast invasion is limited, and interstitial trophoblasts do not invade into the myometrium (106). Furthermore, invasive trophoblasts in rats are intraluminal, replacing the endothelium from within the vessel, while those in mice are perivascular, replacing the endothelium from outside the vessel (106). Vascular changes in rodents and humans are similar. However, the depth of trophoblast invasion into the uterus and uterine vasculature and the intraluminal approach to vessel wall intercalation are most similar between rats and humans. Although the rat may be a more appropriate model of trophoblast invasion, the lack of transgenic rat models has limited investigations of molecular signaling underlying vascular changes. The mechanistic studies described below largely rely on mouse models, for which transgenics are more advanced.

Studies of spiral arteriole remodeling in mice use immunohistochemical staining of endothelial cells with vWF and VSMCs with SMA (107) to identify these key vasculature components. Remodeled spiral arteries have similarities to lymphatic vessels, including reduced SMA, lack of a basement membrane, and a large, dilated lumen (108). Spiral arterioles that were once high-resistance and low-capacitance are thus remodeled to low-resistance high-capacitance vessels that allow for adequate delivery of nutrients and oxygen from the mother to the developing fetus (Figure 3).


The placenta is fully formed by GD10.5 and is composed of three layers, the decidua, junctional zone, and labyrinth, each with unique vasculature. Within the maternal-derived decidua, arterioles undergo angiogenesis and trophoblast invasion after GD8.5 (73). The junctional zone, composed of spongiotrophoblast and glycogen trophoblast cells, is the confluence of maternal and fetal cells of the placenta. The labyrinth portion of the placenta is generally considered the fetal compartment, with fetal vascular endothelium and cytotrophoblast cells in direct contact with maternal blood (109). This type of placentation in rodents and humans, with intimacy of maternal and fetal cells for nutrient exchange, is classified as hemochorial (110).

### Non-trophoblast factors contributing to spiral arteriole remodeling (GD9.0–GD10.5)

Trophoblast-independent rodent spiral arteriole remodeling involves apoptosis and is mediated by angiogenic factors and uNK cells, as described below.

**Apoptosis.** The process of endovascular and interstitial trophoblast invasion requires endothelial cell and VSMC apoptosis, as evidenced by the uterine expression of cleaved caspase-3 (111). Studies using mice deficient in uterine Corin, a serine transmembrane protease necessary for atrial natriuretic peptide (ANP) synthesis, showed that ANP is essential for decidualization and spiral arteriole remodeling (111). ANP-deficient mice demonstrated low expression of uterine TNF-related apoptosis-inducing ligand (TRAIL), which is required for VSMC apoptosis and cyclophilin B–mediated TRAIL receptor upregulation in endothelial cells within spiral arterioles (111). Gene mutations and reduced expression and activity of Corin have been associated with human preeclamptic pregnancy (112), suggesting that these mouse models could help develop a better understanding of the process of spiral arteriole remodeling.

**uNK cells.** uNK cells are the predominant CD45<sup>+</sup> immune cells in the mouse decidua until GD7.5 (73, 113, 114). Following IL-15 stimulation, uNK cells begin secreting key angiogenic and vasoactive factors, including VEGF, PGF, and IFN- $\gamma$ , that are required for vascular remodeling in pregnancy. Retention of decidual vessel SMA in uNK cell-deficient *Rag2<sup>-/-</sup> IL2rg<sup>-/-</sup>* mice results in constricted spiral arterioles at GD9.5 (115). At mid-gestation, mature uNK cells are essential for decidual maintenance. uNK cell-deficient mice have hypocellular decidual basalis at GD9.5–GD10 that becomes hypogranular at GD12 with spiral arterioles resembling those in non-pregnant mice (116).



**Figure 3. Remodeling of spiral arterioles during pregnancy.** Spiral arterioles are lined by endothelial cells and are wrapped on the outside by pericytes and VSMCs. From GD7.5 to GD9.5, the maternal spiral arterioles increase in diameter after embryo-derived trophoblast cells (green) integrate into the lining of the vessel wall. As a result, the maternal vasculature in the pregnant uterus is one of decreased resistance and increased blood flow to nourish the fetus.

*Angiogenic factors.* A balance of pro- and anti-angiogenic factors in the placenta and maternal circulation is required for pregnancy success (Table 4). VEGF interacts directly with maternal uterine endothelial cells to promote angiogenesis and uteroplacental blood flow necessary for fetal circulation by GD9.5. Soluble VEGFR1 (sFlt1) is produced with pathologic placental hypoxia and scavenges free VEGF, preventing adequate placental angiogenesis (117). Paradoxically, VEGF overexpression in the endometrium of mice results in parallel upregulation of anti-angiogenic sFlt1 (118). Ultimately, endometrial VEGF overexpression in early pregnancy results in pathologically enlarged maternal blood spaces within the junctional zone while sparing labyrinth formation before GD11.5 (118).

Although early pregnancy studies are lacking in rats, a transgenic rat model of human preeclampsia was created by breeding of female rats containing a human angiotensinogen transgene with male rats harboring a human renin transgene (119). Uterine arterioles displaying a lower stretch response at GD7.5 correlated with a hypertensive pregnancy resembling the human preeclampsia condition.

#### Trophoblast factors contributing to spiral arteriole remodeling (GD9.0–GD10.5)

Endovascular trophoblast cells replace endothelial cells of the spiral arterioles, transforming them from small resistant vessels to flaccid, large-caliber vessels that are unresponsive to vasoconstrictive agents. This complex process is thought to occur in rodents and women. At the inception, endovascular trophoblast cells form plugs in maternal uterine vessels. This is thought to protect the conceptus from excessively high oxygen levels, and thus placenta formation begins in a hypoxic state (73, 120). Once the plugs dissociate, maternal blood flow to the placenta increases. Notch is involved in trophoblast-dependent spiral arteriole remodeling prior to GD10.5.

*Notch signaling.* Canonical Notch signaling is active in trophoblasts during early pregnancy, and trophoblast subtypes express Notch proteins, including DLL4, Notch2, and Notch4 (121). *Notch2* deletion using the trophoblast-specific driver *Tpbpa-Cre* resulted in inadequate remodeling of spiral arterioles and a decline in placental perfusion at GD10.5 and GD14.5 (122).

#### Eye to the future

There are clear gaps in our understanding of uterine vascular remodeling during early pregnancy. Most studies describe vascular structure, gene expression, and, to some extent, functional changes in rodents (3, 24, 39–42, 57, 123). Very few studies simultaneously investigate signaling pathways and vessel architecture (6, 70). Future studies that combine genetic rodent models and advanced methodologies, such as organoids, organ-on-chip, and omics, with novel imaging techniques will allow integration of gene expression with spatial information and vessel function at different stages of the estrous/menstrual cycle and in early pregnancy.

Endometrial organoids and organ-on-chip represent promising systems that can facilitate study of tissue function in physiologic and pathophysiologic contexts. These systems may provide a suitable approach

**Table 4. GD9.0–GD10.5: factors that regulate angiogenesis and vascular remodeling**

Gene	Method	Expression		Inhibition/overexpression model		Refs.
		Developmental stages	Cell type and/or location	Model	Phenotype	
<i>Anp</i>	Protein (IF)	GD10	uNK cells	Knockout mice	Apoptosis of VSMCs; decreased spiral artery remodeling	135
	Pro-ANP protein (ELISA)	NA	Human endometrial stromal cells	NA	NA	111
<i>Corin</i>	mRNA (RT-qPCR)	GD7.5–GD10.5	Decidual cells	Knockout mice	Reduced trophoblast invasion; smaller arterioles present in decidua at GD12.5	112
<i>Notch2</i>	mRNA (ISH)	GD8.5–GD10.5	Endovascular trophoblast cells	Trophoblast-specific knockout	Reduced caliber of spiral arterioles; reduced placental perfusion	122
<i>Pgf</i>	mRNA (real-time PCR) Protein (IHC)	Weeks 8 and 9 of human gestation	Epithelium and endothelium of the human decidual basalis, and extravillous trophoblasts	NA	NA	136

IF, immunofluorescence; IHC, immunohistochemistry; ISH, in situ hybridization; RT-qPCR, quantitative reverse transcription PCR.

to bridge the gap between 2D cultures and in vivo models (124). Endometrial organoids have been generated using epithelial cells alone or epithelial and stromal cells together but have not yet integrated endothelial cells (125). An implantation-on-chip model has also been constructed to understand the interaction of primary trophoblast cells, which were derived from first-trimester placenta, with uterine stromal and uNK cells. In this model, trophoblast invasion and migration directionality were drastically reduced in the absence of uterine endothelial cells. Further, non-uterine sources of endothelial cells promoted trophoblast invasion to a much lesser extent compared with uterine endothelial cells, supporting the unique properties of organ-specific endothelium (77).

Transcriptomic profiling of the uterine microenvironment with RNA sequencing during different stages of pregnancy is emerging as a powerful tool to determine the expression patterns of angiogenesis genes that are necessary for implantation and pregnancy success (126–128). Single-cell RNA sequencing has the advantage of interrogating the transcriptome of specific cells within the heterogeneous placental tissue composed of maternal- and fetal-derived cells. Information on the cell-to-cell crosstalk between maternal and fetal cells can be investigated more elegantly with spatial transcriptomics (126). Equally important are new imaging advances, such as confocal and light sheet microscopy, that will allow connections between gene expression changes and vascular structural and functional changes during implantation, decidualization, and early pregnancy (125, 129).

## Conclusions

While a lot has been uncovered about uterine vascular changes during early pregnancy, many unanswered questions remain. The principal molecular signals in uterine (maternal) vascular remodeling during early pregnancy and how vessels are modified during clinical conditions such as infertility and miscarriage are not known. The downstream vasoactive and angiogenic pathways directly regulated by ovarian steroid hormones that result in vessel remodeling during the estrous cycle and during pre-implantation stages of pregnancy have not been well studied. Finally, although implied by increased vascular permeability during early pregnancy, how the changes in blood vessel structure relate to function needs further attention. Addressing these gaps in knowledge will provide novel perspectives regarding early pregnancy and allow design of therapeutics to address clinical conditions related to infertility and miscarriage.

## Acknowledgments

RA receives support from March of Dimes grant 5-FY20-209 and NIH R01HD109152; NM receives support from National Institute of Child Health and Human Development T32HD087166; NCD receives support from NIH R01AI148695; and JS receives support from NIH P20GM135002.

Address correspondence to: Rippl Arora, 775 Woodlot Dr., Rm 3312, East Lansing, Michigan 48824, USA. Email: rippl@msu.edu. Or to: Nataki C. Douglas, 185 South Orange Ave, MSB E569, Newark, New Jersey 07103, USA. Email: nd537@njms.rutgers.edu.

1. Ramsey EM. Vascular anatomy. In: Wynn RM, Jollie WP, eds. *Biology of the Uterus*. Springer; 1989:57–68.
2. Greiss FC, Rose JC. Vascular physiology of the nonpregnant uterus. In: Wynn RM, Jollie WP, eds. *Biology of the Uterus*. Springer; 1989:69–87.
3. Rogers PA. Structure and function of endometrial blood vessels. *Hum Reprod Update*. 1996;2(1):57–62.
4. Alsip NL, et al. A new technique for studying the uterine microvasculature in the rat. *Am J Obstet Gynecol*. 1996;175(2):388–395.
5. Takemori K, et al. Scanning electron microscopy study on corrosion cast of rat uterine vasculature during the first half of pregnancy. *J Anat*. 1984;138(pt 1):163–173.
6. Kim M, et al. VEGF-A regulated by progesterone governs uterine angiogenesis and vascular remodelling during pregnancy. *EMBO Mol Med*. 2013;5(9):1415–1430.
7. Mandala M, Osol G. Physiological remodelling of the maternal uterine circulation during pregnancy. *Basic Clin Pharmacol Toxicol*. 2012;110(1):12–18.
8. Osol G, Mandala M. Maternal uterine vascular remodeling during pregnancy. *Physiology (Bethesda)*. 2009;24:58–71.
9. Osol G, et al. Plasticity of the maternal vasculature during pregnancy. *Annu Rev Physiol*. 2019;81:89–111.
10. Ferenczy A. Anatomy and histology of the uterine corpus. In: Kurman RJ, ed. *Blaustein's Pathology of the Female Genital Tract*. Springer; 1994:327–366.
11. Hickey M, Fraser I. Human uterine vascular structures in normal and diseased states. *Microsc Res Tech*. 2003;60(4):377–389.
12. Martínez-Aguilar R, et al. Hypoxia and reproductive health: the presence and role of hypoxia in the endometrium. *Reproduction*. 2021;161(1):F1–F17.
13. Red-Horse K. Lymphatic vessel dynamics in the uterine wall. *Placenta*. 2008;29(suppl a):S55–S59.
14. Girling JE, Rogers PA. The endometrial lymphatic vasculature: function and dysfunction. *Rev Endocr Metab Disord*. 2012;13(4):265–275.
15. Breslin JW, et al. Lymphatic vessel network structure and physiology. *Compr Physiol*. 2018;9(1):207–299.
16. Red-Horse K, et al. Cytotrophoblast induction of arterial apoptosis and lymphangiogenesis in an in vivo model of human placentalization. *J Clin Invest*. 2006;116(10):2643–2652.
17. Gambino LS, et al. Angiogenesis occurs by vessel elongation in proliferative phase human endometrium. *Hum Reprod*. 2002;17(5):1199–1206.
18. Chaudhry R, Chaudhry K. Anatomy, Abdomen and Pelvis: Uterine Arteries. *StatPearls*. StatPearls Publishing; 2022.
19. Lecce G, et al. Presence of estrogen receptor beta in the human endometrium through the cycle: expression in glandular, stromal, and vascular cells. *J Clin Endocrinol Metab*. 2001;86(3):1379–1386.
20. Girling JE, Rogers PA. Recent advances in endometrial angiogenesis research. *Angiogenesis*. 2005;8(2):89–99.
21. Iruela-Arispe ML, et al. Endometrial endothelial cells express estrogen and progesterone receptors and exhibit a tissue specific response to angiogenic growth factors. *Microcirculation*. 1999;6(2):127–140.
22. Critchley HO, et al. Wild-type estrogen receptor (ERbeta1) and the splice variant (ERbeta2) are both expressed within the human endometrium throughout the normal menstrual cycle. *J Clin Endocrinol Metab*. 2002;87(11):5265–5273.
23. Smith SK. Regulation of angiogenesis in the endometrium. *Trends Endocrinol Metab*. 2001;12(4):147–151.
24. Chakraborty I, et al. Differential expression of vascular endothelial growth factor and its receptor mRNAs in the mouse uterus around the time of implantation. *J Endocrinol*. 1995;147(2):339–352.
25. Gargett CE, et al. Focal vascular endothelial growth factor correlates with angiogenesis in human endometrium. Role of intra-vascular neutrophils. *Hum Reprod*. 2001;16(6):1065–1075.
26. Li XF, et al. Angiogenic growth factor messenger ribonucleic acids in uterine natural killer cells. *J Clin Endocrinol Metab*. 2001;86(4):1823–1834.
27. Krikun G, et al. Endometrial angiogenesis: from physiology to pathology. *Ann N Y Acad Sci*. 2004;1034:27–35.
28. Rogers PA, et al. Endometrial microvascular growth in normal and dysfunctional states. *Hum Reprod Update*. 1998;4(5):503–508.
29. Hickey M, et al. Changes in endometrial blood vessels in the endometrium of women with hormone replacement therapy-related irregular bleeding. *Hum Reprod*. 2003;18(5):1100–1106.
30. Maybin JA, Critchley HO. Menstrual physiology: implications for endometrial pathology and beyond. *Hum Reprod Update*. 2015;21(6):748–761.
31. Bertolin K, Murphy BD. Monitoring mouse estrous cycles. In: Croy BA, et al., eds. *The Guide to Investigation of Mouse Pregnancy*. Academic Press; 2013:475–477.
32. Cousins FL, et al. Evidence from a mouse model that epithelial cell migration and mesenchymal-epithelial transition contribute to rapid restoration of uterine tissue integrity during menstruation. *PLoS One*. 2014;9(1):e86378.
33. Shawber CJ, et al. Vascular Notch proteins and Notch signaling in the peri-implantation mouse uterus. *Vasc Cell*. 2015;7:9.
34. Flores D, et al. Mechanical and signaling mechanisms that guide pre-implantation embryo movement. *Development*. 2020;147(24):dev193490.
35. Madhavan MK, et al. Aberrant uterine folding in mice disrupts implantation chamber formation and alignment of

- embryo-uterine axes. *Development*. 2022;149(11):dev200300.
36. Pang SC, et al. The cycling and pregnant mouse: gross anatomy. In: Croy BA, et al., eds. *The Guide to Investigation of Mouse Pregnancy*. Academic Press; 2013:3–19.
  37. Edwards AK, et al. Microscopic anatomy of the pregnant mouse uterus throughout gestation. In: Croy BA, et al., eds. *The Guide to Investigation of Mouse Pregnancy*. Academic Press; 2013:43–67.
  38. Kennedy TG, et al. Prostaglandins and the initiation of blastocyst implantation and decidualization. *Reproduction*. 2007;134(5):635–643.
  39. Goodger AM, Rogers PA. Uterine endothelial cell proliferation before and after embryo implantation in rats. *J Reprod Fertil*. 1993;99(2):451–457.
  40. Goodger AM, Rogers PA. Blood vessel growth and endothelial cell density in rat endometrium. *J Reprod Fertil*. 1995;105(2):259–261.
  41. Rogers PW, et al. Changes in the spatial organization of the uterine vasculature during implantation in the rat. *J Reprod Fertil*. 1982;65(1):211–214.
  42. Tawia SA, Rogers PA. In vivo microscopy of the subepithelial capillary plexus of the endometrium of rats during embryo implantation. *J Reprod Fertil*. 1992;96(2):673–680.
  43. Kennedy TG. Prostaglandins and increased endometrial vascular permeability resulting from the application of artificial stimulus to the uterus of the rat sensitized for the decidual cell reaction. *Biol Reprod*. 1979;20(3):560–566.
  44. Claesson-Welsh L, et al. Permeability of the endothelial barrier: identifying and reconciling controversies. *Trends Mol Med*. 2021;27(4):314–331.
  45. Boshier DP. The pontamine blue reaction in pregnant sheep uteri. *J Reprod Fertil*. 1970;22(3):595–596.
  46. Deanesly R. The role of the fertilized egg: reactions in the guinea-pig uterus at ovo-implantation and after thread traumatization. *J Reprod Fertil*. 1967;14(2):243–248.
  47. Keys JL, et al. Increased uterine vascular permeability at the time of embryonic attachment in the pig. *Biol Reprod*. 1986;34(2):405–411.
  48. Song H, et al. Dysregulation of EGF family of growth factors and COX-2 in the uterus during the preattachment and attachment reactions of the blastocyst with the luminal epithelium correlates with implantation failure in LIF-deficient mice. *Mol Endocrinol*. 2000;14(8):1147–1161.
  49. Stewart CL, et al. Blastocyst implantation depends on maternal expression of leukaemia inhibitory factor. *Nature*. 1992;359(6390):76–79.
  50. Hyder SM, et al. Identification of functional estrogen response elements in the gene coding for the potent angiogenic factor vascular endothelial growth factor. *Cancer Res*. 2000;60(12):3183–3190.
  51. Ma W, et al. Adult tissue angiogenesis: evidence for negative regulation by estrogen in the uterus. *Mol Endocrinol*. 2001;15(11):1983–1992.
  52. Walter LM, et al. Vascular endothelial growth factor-A isoform and (co)receptor expression are differentially regulated by 17beta-oestradiol in the ovariectomised mouse uterus. *Reproduction*. 2010;140(2):331–341.
  53. Hyder SM, Stancel GM. Regulation of angiogenic growth factors in the female reproductive tract by estrogens and progestins. *Mol Endocrinol*. 1999;13(6):806–811.
  54. Hyder SM, et al. Uterine expression of vascular endothelial growth factor is increased by estradiol and tamoxifen. *Cancer Res*. 1996;56(17):3954–3960.
  55. Halder JB, et al. Differential expression of VEGF isoforms and VEGF(164)-specific receptor neuropilin-1 in the mouse uterus suggests a role for VEGF(164) in vascular permeability and angiogenesis during implantation. *Genesis*. 2000;26(3):213–224.
  56. Tan W, et al. Relationship between macrophages in mouse uteri and angiogenesis in endometrium during the peri-implantation period. *Thromb Haemostasis*. 2014;92(7):1021–1027.
  57. Rabbani ML, Rogers PA. Role of vascular endothelial growth factor in endometrial vascular events before implantation in rats. *Reproduction*. 2001;122(1):85–90.
  58. Kennedy TG. Evidence for a role for prostaglandins in the initiation of blastocyst implantation in the rat. *Biol Reprod*. 1977;16(3):286–291.
  59. Kennedy TG, Zamecnik J. The concentration of 6-keto-prostaglandin F1alpha is markedly elevated at the site of blastocyst implantation in the rat. *Prostaglandins*. 1978;16(4):599–605.
  60. Kennedy TG. Evidence for the involvement of prostaglandins throughout the decidual cell reaction in the rat. *Biol Reprod*. 1985;33(1):140–146.
  61. Matsumoto H, et al. Cyclooxygenase-2 differentially directs uterine angiogenesis during implantation in mice. *J Biol Chem*. 2002;277(32):29260–29267.
  62. Lim H, et al. Cyclo-oxygenase-2-derived prostacyclin mediates embryo implantation in the mouse via PPARdelta. *Genes Dev*. 1999;13(12):1561–1574.
  63. Pakrasi PL, Dey SK. Blastocyst is the source of prostaglandins in the implantation site in the rabbit. *Prostaglandins*. 1982;24(1):73–77.
  64. Lau IF, et al. Pregnancy blockade by indomethacin, an inhibitor of prostaglandin synthesis: its reversal by prostaglandins and progesterone in mice. *Prostaglandins*. 1973;4(6):795–803.
  65. Lim H, et al. Multiple female reproductive failures in cyclooxygenase 2-deficient mice. *Cell*. 1997;91(2):197–208.
  66. Sookvanichsilp N, Pulbutr P. Anti-implantation effects of indomethacin and celecoxib in rats. *Contraception*. 2002;65(5):373–378.
  67. Lustgarten Guahmich N, et al. Endothelial deletion of ADAM10, a key regulator of Notch signaling, causes impaired decidualization and reduced fertility in female mice. *Angiogenesis*. 2020;23(3):443–458.
  68. Douglas NC, et al. Vascular endothelial growth factor receptor 2 (VEGFR-2) functions to promote uterine decidual angiogenesis during early pregnancy in the mouse. *Endocrinology*. 2009;150(8):3845–3854.
  69. Marchetto NM, et al. Endothelial Jagged1 antagonizes Dll4/Notch signaling in decidual angiogenesis during early mouse pregnancy. *Int J Mol Sci*. 2020;21(18):6477.
  70. Park YG, et al. Angiotensin-2 regulated by progesterone induces uterine vascular remodeling during pregnancy. *Mol Med Rep*. 2020;22(2):1235–1242.



71. García-Pascual CM, et al. Inhibition of delta-like 4 mediated signaling induces abortion in mice due to deregulation of decidual angiogenesis. *Placenta*. 2014;35(7):501–508.
72. Kuang H, et al. Hormonal regulation of uterine natural killer cells in mouse preimplantation uterus. *J Mol Histol*. 2010;41(1):1–7.
73. Croy BA, et al. Imaging of vascular development in early mouse decidua and its association with leukocytes and trophoblasts. *Biol Reprod*. 2012;87(5):125.
74. Wang C, et al. Granulated metrial gland cells in the murine uterus: localization, kinetics, and the functional role in angiogenesis during pregnancy. *Microsc Res Tech*. 2003;60(4):420–429.
75. Hofmann AP, et al. Uterine natural killer cells pace early development of mouse decidua basalis. *Mol Hum Reprod*. 2014;20(1):66–76.
76. Das A, et al. De novo synthesis of estrogen in pregnant uterus is critical for stromal decidualization and angiogenesis. *Proc Natl Acad Sci U S A*. 2009;106(30):12542–12547.
77. Park JY, et al. A microphysiological model of human trophoblast invasion during implantation. *Nat Commun*. 2022;13(1):1252.
78. Scott CA, et al. Angiopoietin-like gene expression in the mouse uterus during implantation and in response to steroids. *Cell Tissue Res*. 2012;348(1):199–211.
79. Douglas NC, et al. VEGFR-1 blockade disrupts peri-implantation decidual angiogenesis and macrophage recruitment. *Vasc Cell*. 2014;6:16.
80. Sones JL, et al. Angiogenic factor imbalance precedes complement deposition in placentae of the BPH/5 model of preeclampsia. *FASEB J*. 2018;32(5):2574–2586.
81. Sones JL, et al. Genotypic analysis of the female BPH/5 mouse, a model of superimposed preeclampsia. *PLoS One*. 2021;16(7):e0253453.
82. He X, Chen Q. Reduced expressions of connexin 43 and VEGF in the first-trimester tissues from women with recurrent pregnancy loss. *Reprod Biol Endocrinol*. 2016;14(1):46.
83. Croy BA, et al. Uterine natural killer cells: a specialized differentiation regulated by ovarian hormones. *Immunol Rev*. 2006;214:161–185.
84. Li G, et al. Role of uterine natural killer cells in angiogenesis of human decidua of the first-trimester pregnancy. *Sci China C Life Sci*. 2008;51(2):111–119.
85. Chiossone L, et al. In vivo generation of decidual natural killer cells from resident hematopoietic progenitors. *Haematologica*. 2014;99(3):448–457.
86. Degaki KY, et al. Delta-like ligand (DLL)1 expression in early mouse decidua and its localization to uterine natural killer cells. *PLoS One*. 2012;7(12):e52037.
87. Oh MJ, Croy BA. A map of relationships between uterine natural killer cells and progesterone receptor expressing cells during mouse pregnancy. *Placenta*. 2008;29(4):317–323.
88. King A, et al. Evaluation of oestrogen and progesterone receptor expression in uterine mucosal lymphocytes. *Hum Reprod*. 1996;11(5):1079–1082.
89. Borzychowski AM, et al. Functional analysis of murine uterine natural killer cells genetically devoid of oestrogen receptors. *Placenta*. 2003;24(4):403–411.
90. Stewart JA, et al. Endometrial leucocytes: expression of steroid hormone receptors. *J Clin Pathol*. 1998;51(2):121–126.
91. Guo W, et al. Glucocorticoid receptor mediates the effect of progesterone on uterine natural killer cells. *Am J Reprod Immunol*. 2012;67(6):463–473.
92. Zhang J, et al. Ovarian stimulation reduces fetal growth by dysregulating uterine natural killer cells in mice. *Mol Reprod Dev*. 2021;88(9):618–627.
93. Bany BM, et al. Analysis of uterine gene expression in interleukin-15 knockout mice reveals uterine natural killer cells do not play a major role in decidualization and associated angiogenesis. *Reproduction*. 2012;143(3):359–375.
94. Hanna J, et al. Decidual NK cells regulate key developmental processes at the human fetal-maternal interface. *Nat Med*. 2006;12(9):1065–1074.
95. Gibson DA, et al. Estrogen-dependent regulation of human uterine natural killer cells promotes vascular remodelling via secretion of CCL2. *Hum Reprod*. 2015;30(6):1290–1301.
96. Wilkens J, et al. Uterine NK cells regulate endometrial bleeding in women and are suppressed by the progesterone receptor modulator asoprisnil. *J Immunol*. 2013;191(5):2226–2235.
97. Williams AR, et al. The effects of the selective progesterone receptor modulator asoprisnil on the morphology of uterine tissues after 3 months treatment in patients with symptomatic uterine leiomyomata. *Hum Reprod*. 2007;22(6):1696–1704.
98. El-Azzamy H, et al. Dysregulated uterine natural killer cells and vascular remodeling in women with recurrent pregnancy losses. *Am J Reprod Immunol*. 2018;80(4):e13024.
99. Winterhager E, et al. Decidual angiogenesis and placental orientation are altered in mice heterozygous for a dominant loss-of-function Gja1 (connexin43) mutation. *Biol Reprod*. 2013;89(5):111.
100. Plum A, et al. Unique and shared functions of different connexins in mice. *Curr Biol*. 2000;10(18):1083–1091.
101. Laws MJ, et al. Gap junction communication between uterine stromal cells plays a critical role in pregnancy-associated neovascularization and embryo survival. *Development*. 2008;135(15):2659–2668.
102. Nair RR, et al. Reduced expression of gap junction gene connexin 43 in recurrent early pregnancy loss patients. *Placenta*. 2011;32(8):619–621.
103. Yu J, et al. Disruption of gap junctions reduces biomarkers of decidualization and angiogenesis and increases inflammatory mediators in human endometrial stromal cell cultures. *Mol Cell Endocrinol*. 2011;344(1–2):25–34.
104. Watson ED, Cross JC. Development of structures and transport functions in the mouse placenta. *Physiology (bethesda)*. 2005;20:180–193.
105. Simmons DG, et al. Early patterning of the chorion leads to the trilaminar trophoblast cell structure in the placental labyrinth. *Development*. 2008;135(12):2083–2091.
106. Silva JF, Serakides R. Intrauterine trophoblast migration: a comparative view of humans and rodents. *Cell Adh Migr*. 2016;10(1–2):88–110.

107. Zeng L, et al. New insights into the roles of CUL1 in mouse placenta development. *Biochem Biophys Res Commun*. 2021;559:70–77.
108. Pawlak JB, et al. Lymphatic mimicry in maternal endothelial cells promotes placental spiral artery remodeling. *J Clin Invest*. 2019;129(11):4912–4921.
109. Woods L, et al. Regulation of placental development and its impact on fetal growth—new insights from mouse models. *Front Endocrinol (lausanne)*. 2018;9:570.
110. Ain R, et al. Gestation stage-dependent intrauterine trophoblast cell invasion in the rat and mouse: novel endocrine phenotype and regulation. *Dev Biol*. 2003;260(1):176–190.
111. Zhang W, et al. Atrial natriuretic peptide promotes uterine decidualization and a TRAIL-dependent mechanism in spiral artery remodeling. *J Clin Invest*. 2021;131(20):e151053.
112. Cui Y, et al. Role of corin in trophoblast invasion and uterine spiral artery remodelling in pregnancy. *Nature*. 2012;484(7393):246–250.
113. Herington JL, et al. Paracrine signals from the mouse conceptus are not required for the normal progression of decidualization. *Endocrinology*. 2009;150(9):4404–4413.
114. Erlebacher A. Immunology of the maternal-fetal interface. *Annu Rev Immunol*. 2013;31:387–411.
115. Kieckbusch J, et al. MHC-dependent inhibition of uterine NK cells impedes fetal growth and decidual vascular remodelling. *Nat Commun*. 2014;5:3359.
116. Ashkar AA, et al. Assessment of requirements for IL-15 and IFN regulatory factors in uterine NK cell differentiation and function during pregnancy. *J Immunol*. 2003;171(6):2937–2944.
117. Nevo O, et al. Increased expression of sFlt-1 in vivo and in vitro models of human placental hypoxia is mediated by HIF-1. *Am J Physiol Regul Integr Comp Physiol*. 2006;291(4):R1085–R1093.
118. Fan X, et al. Endometrial VEGF induces placental sFLT1 and leads to pregnancy complications. *J Clin Invest*. 2014;124(11):4941–4952.
119. Pulgar VM, et al. Functional changes in the uterine artery precede the hypertensive phenotype in a transgenic model of hypertensive pregnancy. *Am J Physiol Endocrinol Metab*. 2015;309(9):E811–E817.
120. Malassiné A, et al. A comparison of placental development and endocrine functions between the human and mouse model. *Hum Reprod Update*. 2003;9(6):531–539.
121. Levin HI, et al. Dynamic maternal and fetal Notch activity and expression in placentation. *Placenta*. 2017;55:5–12.
122. Hunkapiller NM, et al. A role for Notch signaling in trophoblast endovascular invasion and in the pathogenesis of pre-eclampsia. *Development*. 2011;138(14):2987–2998.
123. Chakraborty I, et al. Developmental expression of the cyclo-oxygenase-1 and cyclo-oxygenase-2 genes in the peri-implantation mouse uterus and their differential regulation by the blastocyst and ovarian steroids. *J Mol Endocrinol*. 1996;16(2):107–122.
124. Song Y, Fazleabas AT. Endometrial organoids: a rising star for research on endometrial development and associated diseases. *Reprod Sci*. 2021;28(6):1626–1636.
125. Yamaguchi M, et al. Three-dimensional understanding of the morphological complexity of the human uterine endometrium. *iScience*. 2021;24(4):102258.
126. Li R, et al. Spatial transcriptomic profiles of mouse uterine microenvironments at pregnancy day 7.5. *Biol Reprod*. 2022;107(2):529–545.
127. Wang W, et al. Single-cell transcriptomic atlas of the human endometrium during the menstrual cycle. *Nat Med*. 2020;26(10):1644–1653.
128. Yang Y, et al. Cell-cell communication at the embryo implantation site of mouse uterus revealed by single-cell analysis. *Int J Mol Sci*. 2021;22(10):5177.
129. Arora R, et al. Insights from imaging the implanting embryo and the uterine environment in three dimensions. *Development*. 2016;143(24):4749–4754.
130. Yu J, et al. Endometrial stromal decidualization responds reversibly to hormone stimulation and withdrawal. *Endocrinology*. 2016;157(6):2432–2446.
131. Sekiguchi K, et al. VEGF receptor 1-expressing macrophages recruited from bone marrow enhances angiogenesis in endometrial tissues. *Sci Rep*. 2019;9(1):7037.
132. Dong H, et al. Effect of acupuncture on endometrial angiogenesis and uterus dendritic cells in COH rats during peri-implantation period. *Evid Based Complement Alternat Med*. 2017;2017:3647080.
133. Wordinger RJ, et al. The immunolocalization of basic fibroblast growth factor in the mouse uterus during the initial stages of embryo implantation. *Growth Factors*. 1994;11(3):175–186.
134. Kennedy TG. Timing of uterine sensitivity for the decidual cell reaction: role of prostaglandins. *Biol Reprod*. 1980;22(3):519–525.
135. Hatta K, et al. Expression of the vasoactive proteins AT1, AT2, and ANP by pregnancy-induced mouse uterine natural killer cells. *Reprod Sci*. 2011;18(4):383–390.
136. Plaisier M, et al. Different degrees of vascularization and their relationship to the expression of vascular endothelial growth factor, placental growth factor, angiopoietins, and their receptors in first-trimester decidual tissues. *Fertil Steril*. 2007;88(1):176–187.